



The Discovery of $\Sigma_b^{(*)}$

Petar Maksimovic, for the CDF collaboration

- Introduction: what is Σ_b ?
- CDF detector, trigger, Λ_b sample
- Blind optimization, background estimates
- Fitting for the $\Sigma_b^{(*)}$ signals
- Systematics, significance



Why $\Sigma_b^{(*)}$?

- Most b-mesons found and their decays studied *extensively*
- Comparatively little is known about heavy baryons
(but several c-baryons recently observed by B factories)
- Finding and studying b-baryons completes and checks the Standard Model
- Measuring masses, decay rates tests theoretical approaches (*description different from B mesons!*)
- Discovering new particles is cool! (And good practice for LHC too)




b-baryons with B=1,C=0,J^P = 1/2⁺, 3/2⁺

have

Notation	Quark content	J ^P	SU(3)	(I,I ₃)	S	B	Mass
Λ_b^0	$b[ud]$	$1/2^+$	3^*	$(0,0)$	0	1	$5619.7 \pm 1.2 \pm 1.2 \text{ MeV}$
Ξ_b^0	$b[su]$	$1/2^+$	3^*	$(1/2,1/2)$	-1	1	5.80 GeV
Ξ_b^-	$b[sd]$	$1/2^+$	3^*	$(1/2,-1/2)$	-1	1	5.80 GeV
Σ_b^+	buu	$1/2^+$	6	$(1,1)$	0	1	5.82 GeV
Σ_b^0	$b\{ud\}$	$1/2^+$	6	$(1,0)$	0	1	5.82 GeV
Σ_b^-	bdd	$1/2^+$	6	$(1,-1)$	0	1	5.82 GeV
$\Xi_b^{0,}$	$b\{su\}$	$1/2^+$	6	$(1/2,1/2)$	-1	1	5.94 GeV
$\Xi_b^{0,}$	$b\{sd\}$	$1/2^+$	6	$(1/2,-1/2)$	-1	1	5.94 GeV
Ω_b^0	bss	$1/2^+$	6	$(0,0)$	-2	1	6.04 GeV
Σ_b^{*+}	buu	$3/2^+$	6	$(1,1)$	0	1	5.84 GeV
Σ_b^{*0}	bud	$3/2^+$	6	$(1,0)$	0	1	5.84 GeV
Σ_b^{*-}	bdd	$3/2^+$	6	$(1,-1)$	0	1	5.84 GeV
Ξ_b^{*0}	bus	$3/2^+$	6	$(1/2,1/2)$	-1	1	5.94 GeV
Ξ_b^{*-}	bds	$3/2^+$	6	$(1/2,-1/2)$	-1	1	5.94 GeV
Ω_b^{*-}	bss	$3/2^+$	6	$(0,0)$	-2	1	6.06 GeV

search
for

The four states of Σ_b





Theoretical expectations

- Predictions from a combinations of potential models, HQET, $1/N_c$ expansion, and lattice

Σ_b property	Expected value (MeV/c ²)
$m(\Sigma_b) - m(\Lambda_b^0)$	180 - 210
$m(\Sigma_b^*) - m(\Sigma_b)$	10 - 40
$m(\Sigma_b^-) - m(\Sigma_b^+)$	5 - 7
$\Gamma(\Sigma_b), \Gamma(\Sigma_b^*)$	$\sim 8, \sim 15$

- Enough as a rough guide for a blind search
- Expect: $\Sigma_b^{(*)}$ is massive enough to decay strongly to $\Lambda_b \pi$, but just barely



Analysis strategy

- Reconstruct Λ_b as:

$$\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$$

$$\Lambda_c^+ \rightarrow p K^- \pi^+$$


- Then combine Λ_b with pions around it to form Σ_b , but treat π^+ and π^- separately:

$$\Sigma_b^{(*)+} \rightarrow \Lambda_b^0 \pi^+$$

$$\Sigma_b^{*-} \rightarrow \Lambda_b^0 \pi^-$$


Tevatron + CDF = *b*-hadron factory

- All species of *b*-hadrons produced!
- Tevatron's has been performing really well: here using $\sim 1.1 \text{ fb}^{-1}$ of data
- CDF has excellent tracking:
 - d_0 resolution
(needed for B physics)
 - p_T resolution
(needed to measure masses)






Reconstructing heavy hadrons




- Decays of hadrons with b and c quarks can be observed with a **Silicon Detector**

- Those CDF can reconstruct are boosted sideways
- Use displacement in transverse plane




Mining b 's from mountains of junk!

- Production rate of b -quarks is very large, but rate of (uninteresting) soft QCD is 1000x larger
- **b -physics program lives and dies by the “trigger system”**
 - very fast electronics
 - examines events in real time
 - decides to keep some events (e.g. those with two displaced tracks)
- *Silicon Vertex Trigger (SVT)* – part of trigger system that finds displaced tracks and triggers on heavy hadrons



Reconstructing Λ_b and Σ_b




- Proton and π from Λ_b usually fire **Two (displaced) Track Trigger** (based on SVT)
- $\bar{B}^0 \rightarrow D^+ \pi^-$ has similar topology, and can be mistaken for $\Lambda_b \rightarrow \Lambda_c^+ \pi^-$ decay
- π from Σ_b comes from primary vertex, *along with tracks from hadronization and Underlying Event*



The largest Λ_b sample in the world

CDF II Preliminary, $L = 1.1 \text{ fb}^{-1}$

$N(\Lambda_b) \sim 3000$





Composition of Λ_b signal window


- 86.4% of Λ_b (all decays)
- 9.3% of B mesons (all decays)
- 4.2% of fake Λ_b (combinatorial)

For Σ_b search, use these numbers to normalize backgrounds on Q distribution

Systematics: shuffle up to 200 events from Λ_b component to two backgrounds

Reconstructing Σ_b

CDF II Preliminary, $L = 1.1 \text{ fb}^{-1}$



“ Λ_b signal region”

“ Λ_b upper sideband”

(source of fake Λ_b background)

- Use Λ_b candidates from “ Λ_b signal region”
- Combine those with prompt tracks to form Σ_b candidates

Reconstructing Σ_b


- Split into two sub-samples:

$\Lambda_b \pi^-$: look for Σ_b^- and Σ_b^{*-}

$\Lambda_b \pi^+$: look for Σ_b^+ and Σ_b^{*+}

- Remove effect of Λ_b resolution by looking at

$$Q \equiv m(\Lambda_b \pi) - m(\Lambda_b) - m_\pi$$






Σ_b optimization

- Use Λ_b signal region
(3σ around Λ_b peak)
- Note: no cut on
 $p_T(\pi$ from Σ_b) !

- Only $\cos\theta^*$ makes substantial difference
- Optimized cuts




Σ_b boost direction
in lab frame

Variable	Cut value
$p_T(\Sigma_b)$	$> 9.5 \text{ GeV/c}$
$ d_0/\sigma_{d_0} $	< 3.0
$\cos\theta^*$	> -0.35




Σ_b optimization: N-1 scan for $\cos\theta^*$

$$\frac{\epsilon(S)}{B} \equiv$$



Backgrounds to worry about





Composition of backgrounds

Background type	Source	Contribution
Λ_b hadronization	PYTHIA	dominant
Combinatorial	Upper Λ_b sideband $m(\Lambda_b) \in [5.8, 7.0]$	small
B meson hadronization	B^0 data	small
All B meson reflections	π_Σ from B hadronization	B^0 PYTHIA
	π_Σ from B decay (D^* , D^{**})	Inclusive b-had MC
	π_Σ from B^{**}	B^0 PYTHIA

Will be ignored from now on



PDF form for background shapes

- All backgrounds modeled with a PDF of this form:

$$f(Q; \alpha, Q_{max}, \gamma) = \left(\frac{Q}{Q_{max}} \right)^{\alpha} e^{-\frac{\alpha}{\gamma} \left(\left(\frac{Q}{Q_{max}} \right)^{\gamma} - 1 \right)}$$



(fits well a whole range of B meson fragmentation shapes)

- Fit separately every background component


(Systematics: try alternative shapes)



Λ_b combinatorial and B hadroniz. bkgs




Λ_b hadronization in PYTHIA



- Need hadronization and Underlying Event background (shape, norm)
- For B mesons, PYTHIA works like a charm
 - cf. SSKT for B_s mixing
- No guarantees for baryons!
- Same as for B mesons, $p_T(\Lambda_b)$ spectrum must be reweighted





Λ_b hadronization: PYTHIA vs data







Reweighting Λ_b hadronization






Λ_b hadronization, after reweighting




A. hadronization background



- Effectively, used PYTHIA to interpolate
- Shape is smooth in Σ_b signal region!

Systematics: use extremes of the track p_T spectrum to reweight

Bkgs before unblinding




- Λ_b hadronization dominates
- Small contribution from
 - B meson bkg
 - Combinatorial
- These backgrounds are fixed when we fit for Σ_b signals




Expected signal (before unblinding)

- Expect 4 peaks:
 - Σ_b^- and Σ_b^{*-} in $\Lambda_b \pi^-$
 - Σ_b^+ and Σ_b^{*+} in $\Lambda_b \pi^+$
- Each peak:
 - Breit-Wigner (\times) Resolution fun.
 - $\Gamma(\Sigma_b)$ predicted by HQET





Detector resolution of measuring Q



- Generated Σ_b PYTHIA MC
- Σ_b states with no natural width
- Checked MC in Σ_c and D^*


Disagreement of 15-20% seen in some cases, use as syst.

$\Gamma(\Sigma_b)$ as a function of M_{Σ_b}

- $\Gamma(\Sigma_b)$ predicted by HQET:
[hep-ph/9406359]

$$\Gamma_{\Sigma_q \rightarrow \Lambda_q \pi} = \frac{1}{6\pi} \frac{M_{\Lambda_q}}{M_{\Sigma_q}} |f_p|^2 |\vec{p}_\pi|^3$$

$$f_p \equiv g_A / f_\pi; g_A = 0.75 \pm 0.05$$




From fit to $\Sigma_c^{(*)++}$ states (use as systematics)


$\Gamma(\Sigma_c^{(*)++})$ in an excellent agreement with PDG



Modeling Σ_b signal peaks




- Natural width from HQET formula
- Dominates over detector resolution!
- Breit-Wigner peaks get wider as $m(\Sigma_b)$ goes up

Candidates per 5 MeV/c²

Box open!

Data	Bkg	excess
416	268	148
406	298	108



The fit

- Backgrounds frozen in the fit
- Signal: 4 peaks, each
 - 2 Breit-Wigners (resolution has 2 Gaussians)
 - $\Gamma(\Sigma_b)$ as a function of center of each peak
- $m(\Sigma_b^*) - m(\Sigma_b)$ common parameter



Fit results

Parameter	Value	MINOS Errors
$\Sigma_b^- Q (\text{MeV}/c^2)$	55.9	(+0.990, -0.959)
Σ_b^- events	60	(+14.8, -13.8)
$\Sigma_b^+ Q (\text{MeV}/c^2)$	48.4	(+2.02, -2.29)
Σ_b^+ events	29	(+12.4, -11.6)
Σ_b^{*-} events	74	(+18.2, -17.4)
Σ_b^{*+} events	74	(+17.2, -16.3)
$\Sigma_b^* - \Sigma_b^- Q (\text{MeV}/c^2)$	21.3	(+2.03, -1.94)

- Only significant correlation between $Q(\Sigma_b^+)$ and $Q(\Sigma_b^*) - Q(\Sigma_b^-)$ (because Σ_b^+ peak is weak...)



Systematics: procedure

- Already listed an array of “variations”:
 - change: Λ_b signal region sample composition, det. resolution, natural width, functional form of background PDFs, extreme reweighting track p_T distribution, etc.
- For each variation:
 - generate 1000 Toy MC experiments with “changed” PDF
 - fit with “baseline” PDF
 - average differences between fit results is the systematic error



Systematics: results

- All small for mass measurements

Parameter	Tracking	Λ_b^0 Comp.	Λ_b^0 Norm.	Λ_b^0 Shape	Reweight	Reso.	Σ_b Width	Total
$Q(\Sigma_b^+)$ (MeV/c ²)	0.06	0.03	0.013	0.013	0.0	0.0	0.01	0.07
	-0.06	0.0	-0.013	0.0	-0.11	-0.014	-0.02	-0.13
$Q(\Sigma_b^-)$ (MeV/c ²)	0.06	0.0	0.009	0.0	0.04	0.0	0.009	0.07
	-0.06	-0.03	-0.002	-0.011	-0.0004	-0.011	-0.005	-0.07
$Q(\Sigma_b^*) - Q(\Sigma_b)$ (MeV/c ²)	0.06	0.05	0.14	0.04	0.32	0.02	0.07	0.37
	-0.06	0.0	-0.13	0.0	0.0	0.0	-0.07	-0.16
Σ_b^+ events	0.0	3.3	2.1	1.2	2.3	0.3	1.8	5.0
	0.0	0.0	-2.1	0.0	-1.8	0.0	-2.0	-3.4
Σ_b^- events	0.0	0.7	2.2	0.3	7.4	0.3	3.4	8.5
	0.0	0.0	-2.2	0.0	0.0	0.0	-3.4	-4.0
Σ_b^{*+} events	0.0	7.3	4.8	2.8	4.6	0.2	0.8	10.3
	0.0	0.0	-4.8	0.0	-2.9	0.0	-0.8	-5.7
Σ_b^{*-} events	0.0	0.4	4.8	0.3	14.7	0.1	1.7	15.6
	0.0	0.0	-4.7	0.0	0.0	0.0	-1.7	-5.0

- Track p_T reweighting largest for yields



Yields (including systematics)

Number of events for each state:

- $N(\Sigma_b^-) = 60^{+14.8}_{-13.8} \text{ (stat)} \quad {}^{+8.4}_{-4.0} \text{ (syst)}$
- $N(\Sigma_b^+) = 29^{+12.4}_{-11.6} \text{ (stat)} \quad {}^{+5.0}_{-3.4} \text{ (syst)}$
- $N(\Sigma_b^{*-}) = 74^{+18.2}_{-17.4} \text{ (stat)} \quad {}^{+15.6}_{-5.0} \text{ (syst)}$
- $N(\Sigma_b^{*+}) = 74^{+17.2}_{-16.3} \text{ (stat)} \quad {}^{+10.3}_{-5.7} \text{ (syst)}$



Significance

- In total, a very significant signal
 - Naïve $S/\sqrt{S+B}$ gives $\sim 9\sigma$
 - P-value calculation $> 5\sigma$: don't have enough Toy MC to probe the 9σ -level (extrapolation too imprecise)
- Strength of signal hypothesis ($4 \Sigma_b$ peaks) best expressed by Likelihood Ratio (LR):

$$LR \equiv \frac{L_{\text{no peak fit}}}{L_{\text{4 peak fit}}} \quad \leftarrow$$

**Evaluate LR
for multiple fit
models and pick the
worst case scenario!**



Likelihood Ratios


- Overall significance
- Four or only two peaks?
- What if one peak is fake?

Hypothesis	$\Delta(-\ln(L))$	$1/LR$
Null	44.7	2.6e19
2 peaks	14.3	1.6e6
No Σ_b^- Peak	10.4	3.3e4
No Σ_b^+ Peak	1.1	3
No Σ_b^{*-} Peak	10.1	2.4e4
No Σ_b^{*+} Peak	9.8	1.8e4


- “It is $\sim 2.6 \times 10^{19}$ more likely that this is a 4 peak Σ_b signal than that it's a background fluctuation!”

Summary

- *Discovered four new particles!*
- ~ 240 events in total
- And measured their masses:
 - $m(\Sigma_b^-) - m(\Lambda_b^0) - m(\pi) = 55.9^{+1.0}_{-1.0}$ (stat) ± 0.1 (syst) MeV/c^2
 - $m(\Sigma_b^+) - m(\Lambda_b^0) - m(\pi) = 48.4^{+2.0}_{-2.3}$ (stat) ± 0.1 (syst) MeV/c^2
 - $m(\Sigma_b^{*-}) - m(\Sigma_b^-) = m(\Sigma_b^{*+}) - m(\Sigma_b^+) = 21.3^{+2.0}_{-1.9}$ (stat) $^{+0.4}_{-0.2}$ (syst) MeV/c^2





Baryons with Up, Down, Strange and Bottom Quarks and Highest Spin ($J = \frac{3}{2}$)





BACKUP SLIDES


Heavy baryon classification



- Σ_b^+ is (uub), decaying to $\Lambda_b^0 \pi^+$
- Σ_b^- is (ddb), decaying to $\Lambda_b^0 \pi^-$

Λ_c and Σ_c states

- Typical decay of Σ -type to Λ -type + π
- For Σ_b , expect similar relationship



Σ_c and Σ_c^*

Λ_c



Tools: Tevatron



- Recently, Tevatron has performed wonderfully
- By now over 2 fb^{-1} delivered to CDF and D0
- This analysis uses 1.1 fb^{-1} delivered to CDF by March 2006



Hopes for the future

- Have about 500 events in $\Lambda_b \rightarrow J/\psi \Lambda$
- Additional 1000 in $\Lambda_b \rightarrow \Lambda_c^+ \pi^-$, but in different triggers
- Potentially another 1k in other channels like $\Lambda_b \rightarrow \Lambda_c 3\pi$

On the shopping list:

- Measure $\Delta m(\Sigma_b)$ in + and – data separately
- Measure production rate relative to Λ_b